

<https://helda.helsinki.fi>

Soil organic matter decomposition and temperature sensitivity after forest fire in permafrost regions in Canada

Aaltonen, Heidi

European Geosciences Union
2017

Aaltonen , H , Palviainen , M , Köster , K , Berninger , F & Pumpanen , J 2017 , Soil organic matter decomposition and temperature sensitivity after forest fire in permafrost regions in Canada . in Geophysical Research Abstracts Vol. 19, EGU General Assembly 2017. . vol. 19 , Geophysical Research Abstracts , vol. 19 , European Geosciences Union , pp. 7359 , European Geosciences Union (EGU) General Assembly 2017 , Wien , Austria , 23/04/2017 .

<http://hdl.handle.net/10138/232481>

cc_by
publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Soil organic matter decomposition and temperature sensitivity after forest fire in permafrost regions in Canada

Heidi Aaltonen (1), Marjo Palviainen (1), Kajar Köster (1), Frank Berninger (1), and Jukka Pumpanen (2)

(1) Department of Forest Sciences, P.O. Box 27, FI-00014 University of Helsinki, Finland, (2) Department of Environmental and Biological Sciences, University of Eastern Finland, P.O. Box 1627, FI-70211 Kuopio, Finland

On the Northern Hemisphere, 24% of soils are underlain by permafrost. These soils contain 50% of the global soil carbon pool. The Northern Hemisphere is also the region which is predicted to be most affected by climate warming and this causes uncertainties over the future of the permafrost. It has been estimated that 25% of permafrost might thaw by 2100, exposing previously frozen carbon pools to decomposition. In addition, global warming is expected to cause increase in the frequency of wild fires, which further increase permafrost melting by removing the insulating organic surface layer. The amount of released soil carbon from permafrost soils after forest fire is affected by degradability and temperature sensitivity of the soil organic matter, as well as soil depth and the stage of succession. Yet the common effect of these factors remains unclear.

We studied how soil respiration and its temperature sensitivity (Q₁₀) vary in different depths and within time by taking soil samples from different fire chronosequence areas (burned 3, 25, 46 and 100 years ago) from permafrost region in Northern Canada (Yukon and Northwest Territories, along Dempster Highway). The samples from three different depths (5, 10 and 30 cm) were incubated in four different temperatures (1, 7, 13 and 19°C) over 24h. Our results showed that the CO₂ fluxes followed the stages of succession, with recently burned sites having lowest rates. The organic matter at 5 cm depth proved to be more labile and temperature sensitive than in deeper depths. The Q₁₀ values, however, did not differ between sites, excluding 30 cm at the most recently burned site that had a significantly higher Q₁₀ value than the other sites. The results implicate that heterotrophic soil respiration decreases on permafrost regions during the first stages after forest fire. At the same time the temperature sensitivity in deeper soil layers may increase.